

Product Name:	Colistin Sodium Methanesulfonate
Product Number:	C073
CAS Number:	8068-28-8
Molecular Formula:	$C_{58}H_{115}N_{16}Na_5O_{28}S_5$
Molecular Weight:	1759.90
Form:	Powder
Appearance:	A white to slightly yellow fine powder
Source:	<i>Bacillus polymyxa</i>
pH:	6.5-8.5
Optical Rotation:	-46° to -51°
Storage Conditions:	2-8 °C
Description:	<p>Colistin sodium methanesulfonate is considered an inactive prodrug of Colistin which means it is inactive until converted into Colistin by cellular enzymes. It is prepared by the reaction of colistin with formaldehyde, then sodium bisulfate. It is less potent and less toxic than colistin sulfate. Colistin targets the bacterial cell membrane, leading to reduced permeability and cell death. It is effective against Gram-negative bacteria. It is freely soluble in aqueous solution.</p> <p>TOKU-E offers three forms of Colistin:</p> <ul style="list-style-type: none">• Colistin sodium methanesulfonate (C073)• <u>Colistin sulfate, USP (C083)</u>• <u>Colistin sulfate, EP (C039)</u> <p>This product is considered a dangerous good. Quantities above 1 g may be subject to additional shipping fees. Please contact us for details.</p>
Mechanism of Action:	Colistin has a bactericidal effect on bacteria by targeting the cell membrane and modifying its permeability.
Spectrum:	Colistin primarily targets and disrupts the cell wall of gram negative bacteria.

Microbiology Applications Colistin is commonly used in clinical *in vitro* microbiological antimicrobial susceptibility tests (panels, discs, and MIC strips) against gram negative microbial isolates. Medical microbiologists use AST results to recommend antibiotic treatment options for infected patients. Representative MIC values include:

- *Shigella* spp. 64 µg/mL -128 µg/mL
- *Haemophilus influenzae* 0.4 µg/mL – 0.8 µg/mL
- For a complete list of colistin MIC values, [click here](#).

References:

Asano T, Kageyama K and Myakumachi (1999) Surface disinfection of resting spores of *Plasmodiophora brassicae* used to infect hairy roots of Brassica spp. *Phytopathol.* 89(4):314-319 PMID 18944777

Bergen, PJ, Li J, Rayner CR, and Nation RL (2006) Colistin methanesulfonate is an inactive prodrug of colistin against *Pseudomonas aeruginosa*. *Antimicrob. Agents Chemother.* 50(6): 1953-1958 PMID 16723551

Falagas ME and Kasiakou SK and Saravolatz LD(2005) Colistin: the revival of polymyxins for the management of multidrug-resistant gram-negative bacterial infections. *Clin. Infect. Dis.* 40(9):1333-1341 PMID 15825037

Koen N, van Breda SV and Loots DT (2018) Elucidating the antimicrobial mechanisms of colistin sulfate on *Mycobacterium tuberculosis* using metabolomics. *Tuberculosis* 1111:14-19 PMID 30029899

Komura S, Kurahashi K (1979) Partial purification and properties of L-2,4-diaminobutyric acid activating enzyme from a polymyxin E producing organism. *J. Biochem (Tokyo)* 86:1013-1021

Leifert C, Ritchie JY and Waites WM (1991) Contaminants of plant-tissue and cell cultures. *World J. Microbiol. Biotechnol.* 7(4):452-469 PMID 24425131

Mueller MJ, Brodschelm W, Spannagl E and Zenk MH (1993) Signaling in the elicitation process is mediated through the octadecanoid pathway leading to jasmonic acid. *Proc. Natl. Acad. Sci. USA* Vol. 90(16):7490-7494 PMID 11607420

Wallace, SJ et al (2012) Interaction of colistin and colistin methanesulfonate with liposomes: colloidal aspects and implications for formulation *J. Pharma Sci* 101 (9):3347-3359 PMID 22623044