



Kanamycin Acid Sulfate, BP PRODUCT DATA SHEET

issue date 01/06/2020

Product Name:	Kanamycin Acid Sulfate, BP
Product Number:	K004
CAS Number:	64013-70-3
Molecular Formula:	$C_{18}H_{36}N_4O_{11} \cdot 2 H_2SO_4$
Molecular Weight:	680.65 g/mol
Form:	Powder
Appearance:	Colorless solid
Solubility:	Water: Freely soluble
Source:	<i>Streptomyces Kanamyceticus</i>
Potency (on a dry basis):	670 IU/mg
Storage Conditions:	2-8 °C
Description:	<p>Kanamycin is an aminoglycoside antibiotic often used to select for bacteria which have been successfully transformed with a plasmid conferring kanamycin resistance. Kanamycin is very soluble in aqueous solution at 92.3 mg/mL.</p> <p>Standard grade kanamycin is composed of a mixture of three different fractions: Kanamycin A, B, and C. TOKU-E offers five forms of kanamycin:</p> <ul style="list-style-type: none">• <u>Kanamycin sulfate</u>• Kanamycin acid sulfate (both BP grade and <u>EP grade</u>)• <u>Kanamycin A sulfate, EvoPure®</u>• <u>Kanamycin B sulfate, EvoPure®</u> <p>EvoPure® products are purified single antibiotic fractions, most >99% pure. High purity EvoPure® kanamycin products can be used to analyze the specific effects of individual kanamycin fractions.</p>
Mechanism of Action:	Aminoglycosides target the 30S ribosomal subunit resulting in an inability to read mRNA ultimately producing a faulty or nonexistent protein.
Spectrum:	Kanamycin is a broad spectrum antibiotic; however, it is mostly used against aerobic gram negative bacteria.

Microbiology Applications Kanamycin acid sulfate is commonly used as a selective agent to select for resistant mammalian, fungal, or bacterial cells that contain the kanMX marker or other kanamycin resistance genes. Kanamycin acid sulfate is typically used at a concentration of 50 µg/mL.

Pryjma, et al. from the University of British Columbia used TOKU-E kanamycin sulfate to select for transformed kanamycin resistant *Campylobacter jejuni* cells: "FdhTU-Modulated Formate Dehydrogenase Expression and Electron Donor Availability Enhance Recovery of *Campylobacter jejuni* following Host Cell Infection"

Media Supplements

Kanamycin can be used as a selective agent in several types of isolation media:

Kanamycin Aesculin Azide Agar - *Enterococci* isolation in food

Perfringens Agar - SFP and TSC selective supplements for the isolation of *Clostridium perfringens*

Plant Biology Applications

Kanamycin is often used in the *Agrobacterium* mediated transformation while using the npt II gene as selection marker. Kaur and Bansal (2010) used kanamycin in combination with cefotaxime to control bacterial growth while transforming tomatoes.

References:

Davis, Bernard D. "Mechanism of Bactericidal Action of Aminoglycosides." *Microbiological Reviews* 51.3 (1987): 341-50.

United States. National Institutes of Health. *Kanamycin Compound Summary*. *PubChem*. Web. 21 Aug. 2012.